


AMENDMENTS TO THE CLAIMS

1. (Currently Amended): Care and/or make-up cosmetic composition comprising a liquid fatty phase comprising at least one volatile silicone oil and at least one volatile non-silicone oil, structured with a gelling system comprising:

1) at least one nylon 611/dimethicone copolymer,
2) at least one non-polymeric organogelling agent, wherein the organogelling agent is selected from the group consisting of:

- (1) - N,N'-bis(dodecanoyl)-1,2-diaminocyclohexane,
- (2) - N,N'-bis(dodecanoyl)-1,3-diaminocyclohexane,
- (3) - N,N'-bis(dodecanoyl)-1,4-diaminocyclohexane,
- (4) - N,N'-bis(dodecanoyl)-1,2-ethylenediamine,
- (5) - N,N'-bis(dodecanoyl)-1-methyl-1,2-ethylenediamine,
- (6) - N,N'-bis(dodecanoyl)-1,3-diaminopropane,
- (7) - N,N'-bis(dodecanoyl)-1,12-diaminododecane,
- (8) - N,N'-bis(dodecanoyl)-3,4-diaminotoluene,
- (9) - at least one compound chosen from the compounds of formula (XV):



in which the groups R⁴⁸, which are identical or different, are chosen from a hydrogen atom and saturated, linear and branched hydrocarbon chains, the said hydrocarbon chains containing from 1 to 6 carbon atoms;

- the groups Z, which are identical or different, each represent a group chosen from the following groups:

-CO-S-R⁴⁹; -CO-NHR⁴⁹; -NH-COR⁴⁹ and -S-COR⁴⁹; in which the groups R⁴⁹, which may be identical or different, are chosen from:

- a hydrogen atom,
- aryl groups,
- aralkyl groups, and
- saturated hydrocarbon chains chosen from linear, branched and cyclic hydrocarbon chains, containing from 1 to 22 carbon atoms, optionally substituted with at least one group chosen from aryl, ester, amide and urethane groups; and/or optionally comprising at least one heteroatom chosen from O, S and N; and/or optionally substituted with at least one fluorine atom and/or one hydroxyl radical,

~~—12-hydroxystearic acid, its salts and its ester or amide derivatives,~~

~~—amides of tricarboxylic acids,~~

~~—esters and amides of N-acylamino acids,~~

~~—diureas of N-acylamino acids,~~


~~—urethane amides of dipeptides,~~

~~—dibenzylidenesorbitol and its derivatives,~~


(10) - sterol derivatives selected from the group consisting of lanosterol, dihydrolanosterol, and cholesterol esters,

~~—cyclo(dipeptides) chosen from cyclo(glycyl-L-alanyl), cyclo(glycyl-L-valyl), cyclo(glycyl-L-leucyl), cyclo(glycyl-L-phenylalanyl), cyclo(L-valyl-L-leucyl), cyclo(L-leucyl-L-leucyl), cyclo(L-phenylalanyl-L-leucyl), cyclo(L-phenylalanyl-L-phenylalanyl), cyclo(L-valyl-L-γ-3,7-dimethyloctylglutamyl), cyclo(L-valyl-L-γ-2-ethylhexylglutamyl), cyclo(L-leucyl-L-γ-ethylglutamyl), cyclo(L-leucyl-L-γ-dodecylglutamyl), cyclo(L-leucyl-L-γ-3,7-dimethyloctylglutamyl), cyclo(L-leucyl-L-γ-benzylglutamyl), cyclo(L-β-butylasparaginyL-phenylalanyl), cyclo(L-γ-dodecylasparaginyL-phenylalanyl), cyclo(L-β-3,7-dimethyloctylasparaginyL-phenylalanyl), cyclo(L-β-2-ethyl~~

~~hexylasparaginyL L-phenylalanyl), ———— cyclo(L-β-3,5,5-~~
~~trimethylhexylasparaginyL L-phenylalanyl) and ———— cyclo(L-β-2-~~
~~ethylbutylasparaginyL L-phenylalanyl),~~
(11) - diamides of formula (XXV) or (XXVI):




or




in which the groups R^{44} , which may be identical or different, represent a saturated, linear or branched C_8 - C_{60} hydrocarbon chain, the group(s) R^{44} optionally comprising a hydroxyl group or at least one heteroatom such as N, O, S or Si, R^{45} is a hydrocarbon-based group chosen from linear, branched and cyclic C_1 to C_{50} groups and C_5 to C_8 arylene groups optionally substituted with one or more C_1 - C_4 alkyl groups, and X represents -O- or -NH-, and mixtures thereof, and

(12) - trans- (1R,2R) -bis(undecylcarbonylamino)cyclohexane of formula:





~~—fluorinated ethers,~~


~~—organogelling agents of formula (XVII):~~






~~in which W and W¹, which may be identical or different, are chosen from CH₂ and CO, and in which Q and Q¹, which may be identical or different, are a hydrocarbon chain chosen from saturated or unsaturated, linear or branched hydrocarbon chains containing at least 6 carbon atoms, and in which s is an integer from 2 to 4;~~

~~—bolaamphiphilic amides derived from amino acids of formulae:~~








in which R^{37} represents:



R^{38} represents:




~~diamide derivatives of benzenedicarboxylic acid and of
valine of formulae:~~




in which L Val represents:



diamides of formula (XXV) or (XXVI):



or



in which the groups R^{44} , which may be identical or different, represent a saturated or unsaturated, linear or branched C_8 - C_{60} hydrocarbon chain, the group(s) R^{44} optionally comprising a hydroxyl group or at least one heteroatom such as N, O, S or Si, R^{45} is a hydrocarbon based group chosen from linear, branched and cyclic C_1 to C_{50} groups and C_5 to C_8 arylene groups optionally substituted with one or more C_1 - C_4

~~alkyl groups, and X represents O or NH, and mixtures thereof,~~

the liquid fatty phase and the gelling system forming a physiologically acceptable medium, and

3) at least one pigment in an amount sufficient to provide a coloring effect to keratin materials upon application.

2-3. (Canceled).

4. (Previously Presented): Composition according to Claim 1, in which the volatile silicone oil has a flash point equal to or greater than 40°C and greater than the softening point of the gelling system.

5. (Previously Presented): Composition according to claim 1, in which the volatile silicone oil is chosen from the group consisting of the following compounds: octyltrimethicone, hexyltrimethicone, octamethylcyclotetrasiloxane D4, dodecamethylcyclohexasiloxane D6, heptamethyloctyltrisiloxane, decamethyltetrasiloxane, dodecamethylpentasiloxane, polydimethylsiloxane of 1.5 cSt, polydimethylsiloxane of 2 cSt, polydimethylsiloxane of 3 cSt, polydimethylsiloxane of 5 cSt, and mixtures thereof.

6. (Previously Presented): Composition according to claim 1, in which the volatile silicone oil has a flash point of 40 to 135°C.

7. (Previously Presented): Composition according to claim 1, in which the liquid fatty phase contains at least 30%

silicone oil with respect to the total weight of the composition.

8. (Previously Presented): Composition according to claim 2, in which the volatile silicone oil represents from 3 to 89.4% of the total weight of the composition.

9. (Previously Presented): Composition according to Claim 1, further comprising at least one filler comprising solid particles.

10. (Original): Composition according to Claim 9, in which the solid particles are hydrophobic particles.

11. (Original): Composition according to Claim 10, in which the solid particles are hydrophilic particles, coated with a film of hydrophobic compound.

12. (Original): Composition according to Claim 9, in which the solid particles are hydrophilic particles and the composition further comprises an amphiphilic silicone.

13. (Previously Presented): Composition according to claim 1, in which the at least one pigment is chosen from zinc oxides, iron oxides, titanium oxides and mixtures thereof.

14-27 (Canceled).

28. (Previously Presented): Composition according to claim 1, in which the at least one nylon 611/dimethicone

copolymer represents from 0.5 to 80% of the total weight of the composition.

29. (Previously Presented): Composition according to claim 1, in which the liquid fatty phase further contains a non-volatile non-silicone oil.

30. (Previously Presented): Composition according to claim 1, in which the liquid fatty phase represents from 5 to 99% of the total weight of the composition.

31. (Canceled):

32. (Canceled):

33. (Canceled):

34. (Canceled):

35. (Canceled):

36. (Canceled):

37. (Previously Presented): Composition according to Claim 1, in which in the at least one volatile non-silicone oil is at least one selected from the group consisting of isododecane, isohexadecane, isohexyl neopentanoate, and isodecyl neopentanoate.

38. (Previously Presented): Composition according to Claim 1, in which in the at least one volatile non-silicone oil is at least one selected from the group consisting of isododecane and isohexadecane.

39. (Previously Presented): Composition according to Claim 38, in which the at least one volatile non-silicone oil is isododecane.

40. (Canceled):

41. (Canceled):


42. (Canceled):

43. (Canceled):

44. (Previously Presented): Composition according to claim 1, in which the organogelling agent comprises at least one compound chosen from:

- N,N'-bis(dodecanoyl)-1,2-diaminocyclohexane,
- N,N'-bis(dodecanoyl)-1,3-diaminocyclohexane,
- N,N'-bis(dodecanoyl)-1,4-diaminocyclohexane,
- N,N'-bis(dodecanoyl)-1,2-ethylenediamine,
- N,N'-bis(dodecanoyl)-1-methyl-1,2-ethylenediamine,
- N,N'-bis(dodecanoyl)-1,3-diaminopropane,
- N,N'-bis(dodecanoyl)-1,12-diaminododecane, and
- N,N'-bis(dodecanoyl)-3,4-diaminotoluene.

45. (Previously Presented): Composition according to claim 1, in which the organogelling agent comprises at least one compound chosen from the compounds of formula (XV):



in which the groups R⁴⁸, which are identical or different, are chosen from a hydrogen atom and saturated, linear and branched hydrocarbon chains, the said hydrocarbon chains containing from 1 to 6 carbon atoms;

- the groups Z, which are identical or different, each represent a group chosen from the following groups: -CO-S-R⁴⁹; -CO-NHR⁴⁹; -NH-COR⁴⁹ and -S-COR⁴⁹; in which the groups R⁴⁹, which may be identical or different, are chosen from:

- a hydrogen atom,
- aryl groups,
- aralkyl groups, and
- saturated hydrocarbon chains chosen from linear, branched and cyclic hydrocarbon chains, containing from 1 to 22 carbon atoms, optionally substituted with at least one group chosen from aryl, ester, amide and urethane groups; and/or optionally comprising at least one heteroatom chosen from O, S and N; and/or optionally substituted with at least one fluorine atom and/or one hydroxyl radical.

46. (Previously Presented): Composition according to Claim 45, in which in the formula (XV), each R⁴⁸ is a hydrogen atom.

47. (Previously Presented): Composition according to claim 45, in which in the formula (XV), each Z is chosen from the groups CONHR⁴⁹ and NH-COR⁴⁹.

48. (Previously Presented): Composition according to claim 45, in which in the formula (XV), R^{49} is chosen from aryl groups; aralkyl groups in which the alkyl portion is a linear or branched alkyl chain comprising 12 to 16 carbon atoms; and linear and branched C_{11} - C_{18} alkyl chains.

49. (Previously Presented): Composition according to claim 45, in which the organogelling agent is chosen from:

- cis-1,3,5-tris(dodecylaminocarbonyl)cyclohexane,
- cis-1,3,5-tris(octadecylaminocarbonyl)cyclohexane,
- cis-1,3,5-tris[N-(3,7-dimethyloctyl)aminocarbonyl]-cyclohexane,
- trans-1,3,5-trimethyl-1,3,5-tris(dodecylaminocarbonyl)cyclohexane, and
- trans-1,3,5-trimethyl-1,3,5-tris(octadecylaminocarbonyl)-cyclohexane.

50-53. (Canceled).

54. (Previously Presented): Composition according to claim 1, in which the organogelling agent is present in a quantity ranging from 0.1% to 80% by weight relative to the total weight of the composition.

55. (Previously Presented): Composition according to claim 1, in which the organogelling agent is present in a quantity ranging from 0.5% to 60% by weight relative to the total weight of the composition.

56. (Previously Presented): Composition according to claim 1, in which the polymer/non-polymeric organogelling agent mass ratio is in the range from 20 to 0.15.

57. (Previously Presented): Composition according to claim 1, wherein the composition further comprises at least one cosmetic or dermatological active agent.

58. (Previously Presented): Composition according to claim 1, wherein the active agent is chosen from essential oils, vitamins, moisturizers, sunscreens, cicatrizing agents and ceramides.

59. (Previously Presented): Composition according to claim 1, wherein comprises at least one additive chosen from dyes that are soluble in polyols or in the fatty phase, antioxidants, essential oils, preserving agents, perfumes, liposoluble polymers, liquid-fatty-phase gelling agents, waxes, gums, resins, surfactants, water, emollients, moisturizers, vitamins, liquid lanolin, essential fatty acids, lipophilic sunscreens or sunscreens that are soluble in polyols, lipid vesicles, and mixtures thereof.

60. (Previously Presented): Composition according to claim 1, wherein the composition further comprises an amphiphilic compound which is liquid at room temperature, having a hydrophilic/lipophilic balance value of less than 12.

61. (Previously Presented): Composition according to claim 1, wherein the composition comprises at least one colouring matter other than a pigment.

62. (Previously Presented): Composition according to claim 1, in the form of an anhydrous stick.

63. (Previously Presented): Composition according to claim 1, in the form of a make-up.

64. (Previously Presented): Composition according to Claim 63, wherein the composition is self-supporting.

65. (Previously Presented): Composition according to claim 1, in the form of a lipstick.

66. (Previously Presented): Composition according to claim 1, in the form of a cake mascara, an eyeliner, a foundation, a lipstick, a blusher, a make-up-removing or deodorant product, a make-up product for the body, an eyeshadow or a face powder, or a concealer product.

67. (Canceled):


68. (Previously Presented): A method of making up a keratinous material comprising applying the composition of claim 1 to the keratinous material.

69. (Currently Amended): A method of making a composition comprising combining at least one nylon 611/dimethicone copolymer

with a liquid continuous fatty phase comprising at least one volatile silicone oil and at least one-volatile non-silicone oil, the liquid fatty phase comprising silicone oil(s) having a flash point equal to or greater than 40°C and greater than the softening point of the and an organogelling agent, wherein the organogelling agent is selected from the group consisting of:

(1) - N,N'-bis(dodecanoyl)-1,2-diaminocyclohexane,

- (2) - N,N'-bis(dodecanoyl)-1,3-diaminocyclohexane,
(3) - N,N'-bis(dodecanoyl)-1,4-diaminocyclohexane,
(4) - N,N'-bis(dodecanoyl)-1,2-ethylenediamine,
(5) - N,N'-bis(dodecanoyl)-1-methyl-1,2-ethylenediamine,
(6) - N,N'-bis(dodecanoyl)-1,3-diaminopropane,
(7) - N,N'-bis(dodecanoyl)-1,12-diaminododecane,
(8) - N,N'-bis(dodecanoyl)-3,4-diaminotoluene,
(9) - at least one compound chosen from the compounds of
formula (XV):



in which the groups R⁴⁸, which are identical or different, are chosen from a hydrogen atom and saturated, linear and branched hydrocarbon chains, the said hydrocarbon chains containing from 1 to 6 carbon atoms;

- the groups Z, which are identical or different, each represent a group chosen from the following groups: -CO-S-R⁴⁹; -CO-NHR⁴⁹; -NH-COR⁴⁹ and -S-COR⁴⁹; in which the groups R⁴⁹, which may be identical or different, are chosen from:

- a hydrogen atom,
- aryl groups,
- aralkyl groups, and
- saturated hydrocarbon chains chosen from linear, branched and cyclic hydrocarbon chains, containing from 1 to 22 carbon atoms, optionally substituted with at least one group chosen from aryl, ester, amide and urethane groups; and/or optionally comprising at least one heteroatom chosen

from O, S and N; and/or optionally substituted with at least one fluorine atom and/or one hydroxyl radical,

~~—12-hydroxystearic acid, its salts and its ester or amide derivatives,~~

~~—amides of tricarboxylic acids,~~

~~—esters and amides of N-acylamino acids,~~

~~—diureas of N-acylamino acids,~~


~~—urethane amides of dipeptides,~~

~~—dibenzylidenesorbitol and its derivatives,~~


(10) - sterol derivatives selected from the group consisting of lanosterol, dihydrolanosterol, and cholesterol esters,

~~—cyclodipeptides chosen from cyclo(glycyl-L-alanyl), cyclo(glycyl-L-valyl), cyclo(glycyl-L-leucyl), cyclo(glycyl-L-phenylalanyl), cyclo(L-valyl-L-leucyl), cyclo(L-leucyl-L-leucyl), cyclo(L-phenylalanyl-L-leucyl), cyclo(L-phenylalanyl-L-phenylalanyl), cyclo(L-valyl-L- γ -3,7-dimethyloctylglutamyl), cyclo(L-valyl-L- γ -2-ethylhexylglutamyl), cyclo(L-leucyl-L- γ -ethylglutamyl), cyclo(L-leucyl-L- γ -dodecylglutamyl), cyclo(L-leucyl-L- γ -3,7-dimethyloctylglutamyl), cyclo(L-leucyl-L- γ -benzylglutamyl), cyclo(L- β -butylasparaginyl-L-phenylalanyl), cyclo(L- γ -dodecylasparaginyl-L-phenylalanyl), cyclo(L- β -3,7-dimethyloctylasparaginyl-L-phenylalanyl), cyclo(L- β -2-ethylhexylasparaginyl-L-phenylalanyl), cyclo(L- β -3,5,5-trimethylhexylasparaginyl-L-phenylalanyl) and cyclo(L- β -2-ethylbutylasparaginyl-L-phenylalanyl),~~

(11) - diamides of formula (XXV) or (XXVI):




or




in which the groups R^{44} , which may be identical or different, represent a saturated, linear or branched C_8 - C_{60} hydrocarbon chain, the group(s) R^{44} optionally comprising a hydroxyl group or at least one heteroatom such as N, O, S or Si, R^{45} is a hydrocarbon-based group chosen from linear, branched and cyclic C_1 to C_{50} groups and C_5 to C_8 arylene groups optionally substituted with one or more C_1 - C_4 alkyl groups, and X represents -O- or -NH-, and mixtures thereof,

(12) - trans-(1R,2R)-bis(undecylcarbonylamino)cyclohexane of formula:



~~fluorinated ethers,~~

~~organogelling agents of formula (XVII):~~




~~in which W and W^1 , which may be identical or different, are chosen from CH_2 and CO, and in which Q and Q^1 , which may be identical or different, are a hydrocarbon chain chosen from saturated or unsaturated, linear or branched~~

Application No. 10/517,390


Response to Office Action dated March 11, 2010

~~hydrocarbon chains containing at least 6 carbon atoms, and in which s is an integer from 2 to 4;~~


~~— bolaamphiphilic amides derived from amino acids of formulae:~~




— 2-alkyl-2-ammoniumisobutyl acetate p-toluenesulphonate salts of formula (XXII):





in which R^{37} represents:



R^{38} represents:




~~diamide derivatives of benzenedicarboxylic acid and of valine of formulae:~~




in which L-Val represents:



diamides of formula (XXV) or (XXVI):



or



in which the groups R^{44} , which may be identical or different, represent a saturated or unsaturated, linear or branched C_8 - C_{60} hydrocarbon chain, the group(s) R^{44} optionally comprising a hydroxyl group or at least one heteroatom such as N, O, S or Si, R^{45} is a hydrocarbon based group chosen from linear, branched and cyclic C_1 to C_{50} groups and C_5 to C_8 arylene groups optionally substituted with one or more C_1 - C_4

~~alkyl groups, and X represents O or NH, and mixtures thereof,~~


and at least one pigment in an amount sufficient to provide a coloring effect to keratin materials upon application, to form a composition which is in the form of a self-supporting solid with a hardness ranging from 20 to 2 000 gf.

70-78. (Canceled).

79. (Previously Presented): Care and/or make-up cosmetic composition comprising a liquid fatty phase comprising at least one volatile silicone oil and at least one volatile non-silicone oil, structured with a gelling system comprising:

1) at least one nylon 611/dimethicone copolymer,
2) at least one non-polymeric organogelling agent, wherein the organogelling agent is selected from the group consisting of:


- N,N'-bis(dodecanoyl)-1,2-diaminocyclohexane,
- N,N'-bis(dodecanoyl)-1,3-diaminocyclohexane,
- N,N'-bis(dodecanoyl)-1,4-diaminocyclohexane,
- N,N'-bis(dodecanoyl)-1,2-ethylenediamine,
- N,N'-bis(dodecanoyl)-1-methyl-1,2-ethylenediamine,
- N,N'-bis(dodecanoyl)-1,3-diaminopropane,
- N,N'-bis(dodecanoyl)-1,12-diaminododecane,
- N,N'-bis(dodecanoyl)-3,4-diaminotoluene,
- at least one compound chosen from the compounds of formula (XV):



in which the groups R⁴⁸, which are identical or different, are chosen from a hydrogen atom and saturated, linear and branched hydrocarbon chains, the said hydrocarbon chains containing from 1 to 6 carbon atoms;

- the groups Z, which are identical or different, each represent a group chosen from the following groups: -CO-S-R⁴⁹; -CO-NHR⁴⁹; -NH-COR⁴⁹ and -S-COR⁴⁹; in which the groups R⁴⁹, which may be identical or different, are chosen from:

- a hydrogen atom,
- aryl groups,
- aralkyl groups, and
- saturated hydrocarbon chains chosen from linear, branched and cyclic hydrocarbon chains, containing from 1 to 22 carbon atoms, optionally substituted with at least one group chosen from aryl, ester, amide and urethane groups; and/or optionally comprising at least one heteroatom chosen from O, S and N; and/or optionally substituted with at least one fluorine atom and/or one hydroxyl radical,
- sterol derivatives selected from the group consisting of lanosterol, dihydrolanosterol, and cholesterol esters, and
- trans-(1R,2R)-bis(undecylcarbonylamino)cyclohexane of formula:




the liquid fatty phase and the gelling system forming a physiologically acceptable medium, and


3) at least one pigment in an amount sufficient to provide a coloring effect to keratin materials upon application.

80. (Previously Presented): Composition according to claim 1, in which the organogelling agent comprises at least one sterol derivative selected from the group consisting of lanosterol, dihydrolanosterol, and cholesterol esters.


81. (Previously Presented): Composition according to claim 1, in which the organogelling agent comprises at least trans-(1R,2R)-bis(undecylcarbonylamino)cyclohexane of formula:



82. (Previously Presented): Composition according to claim 1, in which the organogelling agent comprises at least one compound chosen from diamides of formula (XXV) or (XXVI):



or



in which the groups R^{44} , which may be identical or different, represent a saturated, linear or branched C_8 - C_{60} hydrocarbon chain, the group(s) R^{44} optionally comprising a hydroxyl group or at least one heteroatom such as N, O, S or Si, R^{45} is a hydrocarbon-based group chosen from linear, branched and cyclic C_1 to C_{50} groups and C_5 to C_8 arylene groups optionally substituted with one or more C_1 - C_4 alkyl groups, and X represents -O- or -NH-, and mixtures thereof,